arXiv:0911.0393v1 [math.GT] 2 Nov 2009

WHITNEY'S FORMULAS FOR CURVES ON SURFACES

YURII BURMAN AND MICHAEL POLYAK

Abstract. The classical Whitney formula relates the number of times an oriented plane curve cuts itself to its rotation number and the index of a base point. In this paper we generalize Whitney's formula to curves on an oriented punctured surface $\Sigma_{m,n}$. To define analogs of the rotation number and the index of a base point of a curve γ, we fix an arbitrary vector field on $\Sigma_{m,n}$. Similar formulas are obtained for non-based curves.

1. Introduction

In this paper we study self-intersections of smooth immersed curves on an oriented surface $\Sigma = \Sigma_{m,n}$ of genus m with n punctures. Fix $p \in \Sigma$ and denote $\pi = \pi_1(\Sigma, p)$. We will consider immersions $\gamma : [0,1] \to \Sigma$ with $\gamma(0) = \gamma(1) = p$ and $\gamma'(0) = \gamma'(1)$ — we call them *closed curves with the base point* p. Throughout the paper, all curves are assumed to be generic, i.e. their only singularities are double points of transversal self-intersection, distinct from p.

1.1. **Self-intersections of an immersed curve.** Let γ be a closed curve and d be its self-intersection $d = \gamma(u) = \gamma(v)$, $u < v$. Define $\operatorname{sgn}(d) = +1$ if the orientation of the basis $(\gamma'(u), \gamma'(v))$ coincides with the one prescribed by the orientation of Σ, and $\operatorname{sgn}(d) = -1$ otherwise, see Figure 1a. Turaev [4] constructed an important element of the group ring $\mathbb{Z}[\pi]$ corresponding to γ, in the following way. Let $\tau_d(\gamma) \in \pi$ be the homotopy class of a loop $\gamma(t)$ with $t \in [0,u] \cup [v,1]$. Denote $D(\gamma)$ the set of double points of γ and define the element $\langle\gamma\rangle \in \mathbb{Z}[\pi]$ by

$$\langle\gamma\rangle = \sum_{d \in D(\gamma)} \operatorname{sgn}(d)\tau_d(\gamma). \tag{1}$$

In particular, for a curve on $\Sigma = \mathbb{R}^2$ one has $\mathbb{Z}[\pi] = \mathbb{Z}$ so $\langle\gamma\rangle = \sum_{d \in D(\gamma)} \operatorname{sgn}(d) \in \mathbb{Z}$.

Example 1. Let $\Sigma = \Sigma_{0,2} = \mathbb{R}^2 \setminus \{0\}$ and denote by g the generator of π (represented by a small loop around 0). For a curve γ shown in Figure 2a we have $\langle\gamma\rangle = g^2 - g$; signs of self-intersections and the corresponding curves τ_d are shown in Figure 2b.

For a curve γ on $\Sigma = \mathbb{R}^2$ one may define its *Whitney index* (or winding number) $\mathrm{w}(\gamma)$ as the number of full rotations made by the tangent vector $\gamma'(t)$ around the origin, as t moves from 0 to 1.

For a curve γ on $\Sigma = \mathbb{R}^2$ and $x \in \mathbb{R}^2 \setminus \gamma$ one can also define $\operatorname{ind}(\gamma, x)$ as the number of times the curve γ circles around x. In other words, it is the linking

2000 *Mathematics Subject Classification.* 57N35, 57R42, 57M20.

Key words and phrases. Whitney formula, curves on surfaces, rotation number, self-intersections.

The first author was partially supported by the ISF grant 1261/05.

number of a 1-cycle $[\gamma]$ with the 0-chain $[\infty]-[x]$ (composed of a point near infinity taken with the positive sign and x taken with the negative sign). It can be calculated as the intersection number of the curve γ with any ray starting in x and going to infinity. If γ is a plane curve with a base point p, we define $\operatorname{ind}(\gamma, p) \in \frac{1}{2}\mathbb{Z}$ by averaging the values of ind on two components of $\mathbb{R}^2 \setminus \gamma$ adjacent to p.

H. Whitney in [5] considered closed curves with a base point on $\Sigma = \mathbb{R}^2$ and showed that

Theorem 2 ([5])**.** *Let $\gamma : [0,1] \to \mathbb{R}^2$ be a generic immersed curve with the base point $p = \gamma(0) = \gamma(1)$. Then*

$$\langle \gamma \rangle = -\operatorname{w}(\gamma) + 2\operatorname{ind}(\gamma, p). \tag{2}$$



FIGURE 1. Signs of self-crossings and some simple curves on $\mathbb{R}^2$.

Example 3. For a curve γ shown on Figure 1b we have $\langle \gamma \rangle = +1$, $\operatorname{w}(\gamma) = 2$, and $\operatorname{ind}(\gamma, p) = 3/2$. For a curve γ shown on Figure 1c we have $\langle \gamma \rangle = +1 - 1 = 0$, $\operatorname{w}(\gamma) = 1$, and $\operatorname{ind}(\gamma, p) = 1/2$.

1.2. **Curves on surfaces.** The main results of this paper are generalizations of Theorem 2 for curves on surfaces. We define appropriate surface versions of expressions in both sides of (2) and relate them.

To define an analog of the Whitney index for a curve $\gamma : S^1 \to \Sigma$, we fix a vector field X on Σ having no zeros on the curve γ. Let $w(\gamma, X)$ be the number of rotations of $\gamma'(t)$ relative to X. It can be calculated as the algebraic number of points in which $\gamma'(t)$ looks in the direction of X; each such point is counted with a positive sign, if γ' turns counter-clockwise relative to X in a neighborhood of $\gamma(t)$ and with a negative sign otherwise. It is clear that $w(\gamma, X)$ does not change under the homotopies of γ and X (as long as γ stays an immersion and X has no zeros on its image). Define the *Whitney index* of γ as $\operatorname{w}(\gamma, X) \stackrel{\text{def}}{=} w(\gamma, X)[\gamma] \in \mathbb{Z}[\pi]$ where $[\gamma]$ is the class of homotopy represented by the curve γ.

In particular, if $\Sigma = \Sigma_{0,1} = \mathbb{R}^2$ and $X = \frac{\partial}{\partial x}$ is a horizontal vector field directed from left to right, then π is trivial and $\operatorname{w}(\gamma, X) = \operatorname{w}(\gamma)$ is the usual Whitney index of γ.

Example 4. Let's return to Example 1. Let $X = f(x,y)(x\frac{\partial}{\partial x} + y\frac{\partial}{\partial y})$ be a radial vector field on $\Sigma = \Sigma_{0,2} = \mathbb{R}^2 \setminus \{0\}$; the function $f(x,y) > 0$ is chosen so that the field X is smooth. One may check that $\operatorname{w}(\gamma, X) = (\operatorname{w}(\gamma) - k)g^k$, where $\operatorname{w}(\gamma)$ is the usual Whitney index of γ, k is the number of times γ circles around $\{0\}$, and



FIGURE 2. A curve on a punctured plane and a radial vector field.

g is the generator of $\pi = \pi_1(\mathbb{R}^2 \smallsetminus \{0\})$. In particular, for the curve γ shown on Figure 2a, the number of rotations of γ' relative to X is -1 (indeed, there is only one tangency point of X with γ, denoted by a in Figure 2c; its sign is -1). Also, $[\gamma] = g^2$, thus $\mathrm{w}(\gamma, X) = -g^2$.

The results obtained in this article generalize those of the papers [2] and [1] where the curves on $\Sigma_{0,1} = \mathbb{R}^2$ and $\Sigma_{1,0} = \mathbb{T}^2$ were considered; the methods we use are similar to those of [1].

We consider both curves with a base point (Theorem 5), and curves without base points (Theorem 9). Both theorems are proved by similar methods, so we collected all proofs in Section 4.

2. Curves with base points

2.1. **Making loops from pieces.** For two generic paths $\gamma_i : [a_i, b_i] \to \Sigma$, $i = 1, 2$ with $\gamma_1(a_1) = \gamma_2(b_2) = p$ we may define an element similar to (1) as follows. Let $d = \gamma_1(u) = \gamma_2(v)$ be an intersection point of γ_1 and γ_2, with $u \neq a_1, b_1$, $v \neq a_2, b_2$. Again, define $\mathrm{sgn}(d) = +1$ if the orientation of the basis $(\gamma_1'(u), \gamma_2'(v))$ coincides with the one prescribed by the orientation of Σ, and $\mathrm{sgn}(d) = -1$ otherwise. Let $\tau_d(\gamma_1, \gamma_2) \in \pi$ be the homotopy class of a loop composed of two arcs: $\gamma_1(t_1)$ with $t_1 \in [a_1, u]$, and $\gamma_2(t_2)$ with $t_2 \in [v, b_2]$, see Figure 3a.



FIGURE 3. Making a loop out of two and three pieces.

Denote by $D(\gamma_1, \gamma_2)$ the set of intersections of γ_1 and γ_2 and define

$$\langle \gamma_1, \gamma_2 \rangle = \sum_{d \in D(\gamma_1, \gamma_2)} \mathrm{sgn}(d) \tau_d(\gamma_1, \gamma_2). \tag{3}$$

2.2. **Intersections and τ-indices.** Let X be a vector field on the surface Σ. Throughout the paper, we will assume that every trajectory of X is infinitely extendable, so a one-parametrical group Φ_t of diffeomorphisms generated by X is well-defined. Denote by $\Phi(a)$ its integral trajectory starting at the point $a \in \Sigma$.

Given X, we generalize formula (1) as follows. Let $\gamma : [0,1] \to \Sigma$ be a generic immersed curve with the base point $p = \gamma(0) = \gamma(1)$. Fix $T > 0$ such that $\Phi_{\pm T}(p) \notin \gamma$. Denote by Φ_- and Φ_+ the negative (resp. positive) T-time semi-trajectories $\Phi_t(p)$, $t \in [-T, 0]$ (resp. $t \in [0, T]$) of the base point p.

Define the *T-time index of p with respect to γ* by

$$\operatorname{ind}_T(\gamma, p) = \frac{1}{2}\big(\langle \gamma, \Phi_- \rangle + \langle \Phi_+, \gamma \rangle\big) \in \frac{1}{2}\mathbb{Z}[\pi] \tag{4}$$

Denote by $\gamma_T : [0,1] \to \Sigma$ a T-shift of γ along X: $\gamma_T(t) = \Phi_T(\gamma(t))$. Suppose that X does not vanish on γ and all the intersections of γ_T with γ are transversal double points $d = \gamma_T(u) = \gamma(v)$. Define $\operatorname{sgn}(d) = +1$ if the orientation of the basis $(\gamma_T'(u), \gamma'(v))$ coincides with the orientation of Σ, and $\operatorname{sgn}(d) = -1$ otherwise. Let $\tau_{d,T}(\gamma) \in \pi$ be the homotopy class of a loop composed of three arcs: $\gamma(t_1)$ with $t_1 \in [0, u]$, $\Phi_{t_2}(\gamma(u))$ with $t_2 \in [0, T]$, and $\gamma(t_3)$ with $t_3 \in [v, 1]$, see Figure 3b.

Between all such intersection points d pick only the ones with $u < v$; denote this set by $D_T(\gamma)$. Define the element $\langle \gamma \rangle_T \in \mathbb{Z}[\pi]$ by

$$\langle \gamma \rangle_T = \sum_{d \in D_T(\gamma)} \operatorname{sgn}(d) \tau_{d,T}(\gamma). \tag{5}$$

In particular, for a curve on $\Sigma = \mathbb{R}^2$ one has $\mathbb{Z}[\pi] = \mathbb{Z}$ so $\langle \gamma \rangle_T \in \mathbb{Z}$.

For a small $T = \varepsilon$ intersections $\gamma(u) = \gamma_\varepsilon(v)$ appear near double points of γ and near points in which γ is tangent to X (the condition $u < v$ means in this case that the tangent vector should be directed in the direction of X). After checking the signs, we see that the contribution of double points to $\langle \gamma, \gamma_\varepsilon \rangle$ equals $\langle \gamma \rangle$, while the contribution of points in which γ is tangent to X equals $\mathrm{w}(\gamma, X)$ (recall that we consider $\mathrm{w}(\gamma, X)$ as a multiple of the class $[\gamma]$) and thus obtain

$$\langle \gamma \rangle_\varepsilon = \langle \gamma \rangle + \mathrm{w}(\gamma, X) \tag{6}$$

Theorem 5. *Let $\gamma : [0,1] \to \Sigma$ be a generic immersed curve with the base point $p = \gamma(0) = \gamma(1)$. Let X be a vector field which does not vanish on γ and is transversal to γ at the base point p. Suppose that $T > 0$ is such that all intersections of γ with γ_T are transversal double points distinct from p. Then*

$$\langle \gamma \rangle = \langle \gamma \rangle_T - \mathrm{w}(\gamma, X) + 2 \operatorname{ind}_T(\gamma, p). \tag{7}$$

Remark. For a small $T = \varepsilon$ we have $\operatorname{ind}_\varepsilon(\gamma, p) = 0$ and $\langle \gamma \rangle_\varepsilon$ is given by (6), so in this case equality (7) is trivial.

Example 6. Let $\Sigma = \Sigma_{0,1} = \mathbb{R}^2$ and $X = \frac{\partial}{\partial x}$ be a horizontal vector field directed from left to right. Here π is trivial, and $\mathrm{w}(\gamma, X) = \mathrm{w}(\gamma)$ is the usual Whitney index of γ. For a large T, γ_T is shifted far from γ, so $\langle \gamma \rangle_T = 0$. Also, $\operatorname{ind}_T(\gamma, p)$ counts intersections of γ with both horizontal rays emanating from p to the left and to the right. So it equals to the index $\operatorname{ind}(\gamma, p)$ of the base point, introduced in Section 1.1. Equation (7) turns in this case into the classical Whitney's formula (2).

Example 7. Let's return to Example 4. For a large T the curve γ_T is shifted far from γ, so $\langle \gamma \rangle_T = 0$. The curve γ shown in Figure 2a has no intersections with Φ_+, and only one intersection with Φ_-, denoted by b in Figure 2c; its sign is -1.

The corresponding curve τ_b is depicted there in bold. It represents a class g, thus $2\operatorname{ind}_T(\gamma, p) = -g$. This agrees with $\langle\gamma\rangle + \mathrm{w}(\gamma, X) = (g^2 - g) - g^2 = -g$.

2.3. **Formula for an infinite time shift.** Let us consider the behavior of formula (7) when $T \to \infty$.

Proposition 8. *Let $\gamma : [0, 1] \to \Sigma$ be a generic immersed curve with the base point $p = \gamma(0) = \gamma(1)$. Let X be a vector field which does not vanish on γ. Suppose that the trajectory $\Phi(p)$ of the base point intersects γ only in a finite number of points. Then limits $\operatorname{ind}(\gamma, p) = \lim\limits_{T\to\infty} \operatorname{ind}_T(\gamma, p)$ and $\langle\gamma\rangle_\infty = \lim\limits_{T\to\infty} \langle\gamma\rangle_T$ are well defined and satisfy*

$$\langle\gamma\rangle = \langle\gamma\rangle_\infty - \mathrm{w}(\gamma, X) + 2\operatorname{ind}(\gamma, p). \tag{8}$$

Proof. By the assumption, $\Phi(p)$ intersects γ in a finite number of points $\Phi_{t_i}(p)$, $i = 1, 2, \dots, k$ with $t_1 < t_2 < \cdots < t_k$. Thus when $T > \max(-t_1, t_k)$, the index $\operatorname{ind}_T(\gamma, p)$ does not change and $\lim_{T\to\infty} \operatorname{ind}_T(\gamma, p)$ is well-defined.

Now the statement follows from Theorem 5. Indeed, note that in addition to $\operatorname{ind}_T(\gamma, p)$, only one term in equation (7) depend on T, namely $\langle\gamma\rangle_T$. Since equation (7) is satisfied for any T, we conclude that $\langle\gamma\rangle_T$ also does not change when $T > \max(-t_1, t_k)$. Therefore, $\lim_{T\to\infty}\langle\gamma\rangle_T$ is also well-defined. Since the equality (7) holds for any T, it is satisfied also in the limit $T \to \infty$. □

Remark. One can calculate $\langle\gamma\rangle_T$ when $T \to \infty$ in another way, using intersections with the graph of the so-called separatrices of X. Separatrices of X are trajectories of X which bound regions of different dynamics of the flow on Σ. The idea is that, under appropriate assumptions on X, as $T \to \infty$, the curve γ_T is attracted into a neighborhood of the union of separatrices of X.

An interested reader may easily visualize it in the simplest case of a gradient vector field of a Morse function on $\Sigma = \Sigma_{m,0}$. In this case, separatrices form a graph with vertices in the critical points, with each pair of neighboring critical points connected by two edges (resulting in a total of $2m + 2$ vertices and $4m + 4$ edges).

For a general vector field this approach involves a number of technicalities and requires a lengthy treatment, so we decided to omit it.

3. Curves without base points

Our goal is to repeat all constructions of Section 2 for curves without base points. In all formulas we will be using now free loops instead of based loops; denote by Ω the loop space of Σ. Let $\gamma : S^1 \to \Sigma$ be an oriented curve without a base point; we are to define an appropriate analogue of $\langle\gamma\rangle$ in this case. Let d be a self-intersection of γ. Smoothing γ in d with respect to the orientation, we obtain two closed curves: γ_d^l and γ_d^r, where the tangent vector of γ_d^l rotates clockwise in the neighborhood of d, and that of γ_d^r rotates counter-clockwise. See Figure 4a. Denote $D(\gamma)$ the set of double points of γ and define $\langle g\rangle \in \mathbb{Z}[\Omega]$ by

$$\langle\gamma\rangle = \sum_{d\in D(\gamma)} ([\gamma_d^l] - [\gamma_d^r]).$$

Define the Whitney index in the non-based case as $\mathrm{w}(\gamma, X) \stackrel{\text{def}}{=} w(\gamma, X)([\gamma] - 1) \in \mathbb{Z}[\Omega]$ where 1 is the class of the trivial loop, and $w(\gamma, X)$ is, like for pointed curves,



FIGURE 4. Smoothing a crossing and constructing loops from pieces

the number of full rotations of $\gamma'(t)$ relative X. (Note that the definition of $w(\gamma, X)$ does not require the base point.)

The definition of $\langle\gamma\rangle_T$ remains almost the same as in Section 2.2, with few straightforward changes. Namely, we drop the requirement that $u < v$ when we consider intersection points $d = \gamma(v) = \gamma_T(u)$ of γ with γ_T, and the corresponding loop $\tau_{d,T}(\gamma)$ consists of just two arcs: a piece of γ parameterized by the arc $\overline{vu}$ of the circle, and $\Phi_t(\gamma(u))$ for $0 \le t \le T$. See Figure 4b (the loop $\tau_{d,T}(\gamma)$ is shown in bold). The following theorem is an analogue of Theorem 5 for curves without base points:

Theorem 9. *Let $\gamma : S^1 \to \Sigma$ be a generic immersed curve without the base point. Let X be a vector field on Σ which does not vanish on γ. Suppose that $T > 0$ is such that all intersections of γ with γ_T are transversal double points. Then*

$$\langle\gamma\rangle = \langle\gamma\rangle_T - \mathrm{w}(\gamma, X). \tag{9}$$

Thus $\langle\gamma\rangle$ provides a simple obstruction for pushing a curve off itself:

Corollary 10. *Let γ be a generic curve on Σ. If $\langle\gamma\rangle \notin \mathbb{Z}[[\gamma] - 1] \subset \mathbb{Z}[\Omega]$ then the curve γ cannot be pushed off itself by a flow of vector field, i.e. any shifted copy γ_T intersects the initial curve γ.*

Example 11. Let $\Sigma = \Sigma_{0,3}$ be a doubly-punctured plane and γ be a figure-eight curve with lobes going around the punctures of Σ. Then γ cannot be pushed off itself.

Note also that the only term in formula (9), which depends on T, is $\langle\gamma\rangle_T$. Thus it is, in fact, independent of T and different values of T give the same value of $\langle\gamma\rangle_T$.

4. Proofs of Theorems 5 and 9

4.1. **Idea of the proofs.** The main idea is rather simple. We interpret both sides of the formula (7) as two different ways to compute an intersection number of two 2-chains in a 4-manifold. Very roughly speaking, the manifold is $\Sigma \times \Sigma$ and two chains are $\{(\gamma(u), \gamma(v)) \mid \ 0 \le u \le v \le 1\}$ and the diagonal $\{(x, x) \mid x \in \Sigma\}$. Their intersection is the left part of the formula (the sum of signs of double points). The same intersection number can be also computed by intersecting the boundary of the first chain with a 3-chain, constructed by homotopy $\{(x, \Phi_t(x)) \mid \ 0 \le t \le T\}$ of the diagonal. This comprises the right hand side of the formula. The reality is somewhat more complicated so some technicalities are involved. In particular,

to split the formula by homotopy classes $\tau \in \pi$ we have to work in the universal covering space $\widetilde{\Sigma}$ of Σ. Also, to push the boundary of the first chain off the diagonal, we take a certain ε cut-off. Theorem 9 is completely similar, except that we use a slightly different configuration space. We treat the case of Theorem 5 in details and indicate modifications needed for Theorem 5.

4.2. **Configuration spaces.** Denote by $\widetilde{\Sigma}$ the set of homotopy classes of paths $\xi : [a, b] \to \Sigma$ such that $\xi(a) = p$; $\widetilde{\Sigma}$ is a universal covering space for Σ and inherits its orientation. The projection $\mathrm{proj} : \widetilde{\Sigma} \to \Sigma$ maps a path ξ into its final point $\xi(b)$. The vector field X can be lifted to the vector field $\widetilde{X}$ on $\widetilde{\Sigma}$ such that $\mathrm{proj}(\widetilde{\Phi}(a)) = \Phi(\mathrm{proj}(a))$, where $\widetilde{\Phi}$ means the trajectory of $\widetilde{X}$.

Consider a configuration space $C = \{(u, v) \mid \ 0 < u < v < 1\}$ of (ordered) pairs of points on $[0, 1]$. To compactify C, we pick a small $\varepsilon > 0$ and take an ε-cut-off:

$$C^\varepsilon = \{(u, v) \mid \ \varepsilon \le u < u + \varepsilon \le v \le 1 - \varepsilon\} \subset C.$$

Note that apart from compactifying C, the condition $u + \varepsilon \le v$ used in this cut-off allows us to push the boundary of C^ε off the diagonal $u = v$. Topologically, C^ε is a closed 2-simplex, whose boundary consists of three intervals, on which $u = \varepsilon$, $v = 1 - \varepsilon$, and $u + \varepsilon = v$, respectively.

The curve γ defines an evaluation map $\mathrm{ev} : C \to \Sigma \times \Sigma$, $\mathrm{ev}(u, v) \stackrel{\mathrm{def}}{=} (\gamma(u), \gamma(v))$ — we forget the curve itself and leave only the two marked points. For a fixed class $\tau \in \pi$ define $\mathrm{ev}_\tau : C^\varepsilon \to \widetilde{\Sigma} \times \widetilde{\Sigma}$ by

$$\mathrm{ev}_\tau : (u, v) \mapsto (\xi_1, \xi_2)$$

where ξ_1 is an arc $\gamma(t)$ with $t \in [0, u]$ (so its final point is $\gamma(u)$) and ξ_2 is made of the loop τ followed by an arc $\gamma(1 - t)$ with $t \in [0, 1 - v]$ (so its final point is $\gamma(v)$), see Figure 5. From the definition of ev_τ we immediately get $\mathrm{proj} \circ \mathrm{ev}_\tau(u, v) = \mathrm{ev}(u, v)$, so ev_τ is a lift of the evaluation map.



FIGURE 5. Lifting the evaluation map

The diagonal

$$\Delta = \{(x, x) \mid x \in \widetilde{\Sigma}\}$$

is a proper 2-dimensional submanifold of $\widetilde{\Sigma} \times \widetilde{\Sigma}$, which inherits the orientation of $\widetilde{\Sigma}$ (as the image of the map $x \mapsto (x, x)$). For a sufficiently small ε the intersection number $\mathrm{I}(\mathrm{ev}_\tau(C^\varepsilon), \Delta)$ of the image of C^ε with Δ is well-defined and independent of ε, since $\mathrm{ev}_\tau(C^\varepsilon)$ is oriented, compact, and its boundary does not intersect Δ.

Denote by $\langle\gamma\rangle^\tau$ the coefficient of τ in $\langle\gamma\rangle$, see (1).

Lemma 12. *We have* $\mathrm{I}(\mathrm{ev}_\tau(C^\varepsilon), \Delta) = \langle\gamma\rangle^\tau$

Proof. Indeed, a pair (u,v) gives an intersection point of $\mathrm{ev}_\tau(C^\varepsilon)$ with Δ iff the homotopy classes of paths ξ_1 and ξ_2 coincide, i.e., endpoints $\gamma(u)$ and $\gamma(v)$ of these paths coincide and the homotopy class of $\xi_2^{-1}\circ\xi_1$ is trivial. This means that $d=\gamma(u)=\gamma(v)$ is a self-intersection point of γ and the homotopy class of the loop $\gamma(t)$ with $t\in[0,u]\cup[v,1]$ equals to τ. The local intersection sign is easy to compute and a direct check assures that orientations of $\mathrm{ev}_\tau(C^\varepsilon)$ and Δ give the positive orientation of $\widetilde{\Sigma}\times\widetilde{\Sigma}$ iff $\mathrm{sgn}(d)=+1$. □

Consider the 2-chain $\Delta_T \stackrel{\text{def}}{=} \{(x,\widetilde{\Phi}_T(x)) \mid x\in\widetilde{\Sigma}\}$. The homotopy $W_T=\{(x,\widetilde{\Phi}_t(x)) \mid 0\le t\le T\}$ is a 3-chain such that $\partial W_T=\Delta_T-\Delta$. Denote I the intersection number, so that one has

$$\mathrm{I}(\mathrm{ev}_\tau(C^\varepsilon),\Delta)=\mathrm{I}(\mathrm{ev}_\tau(C^\varepsilon),\Delta_T)-\mathrm{I}(\mathrm{ev}_\tau(\partial C^\varepsilon),W_T) \tag{10}$$

Denote by $\langle\gamma\rangle_T^\tau$ the coefficient of τ in $\langle\gamma\rangle_T$, see (5).

Lemma 13. *We have* $\mathrm{I}(\mathrm{ev}_\tau(C^\varepsilon),\Delta_T)=\langle\gamma\rangle_T^\tau$

The proof repeats the proof of Lemma 12 above.

In view of (10) it remains to compute $\mathrm{I}(\mathrm{ev}_\tau(\partial C^\varepsilon),W_T)$. Each intersection point of $\mathrm{ev}_\tau(\partial C^\varepsilon)$ with W_T corresponds to a pair $(u,v)\in\partial C^\varepsilon$, such that the endpoint of the corresponding path ξ_2 is obtained from the endpoint of ξ_1 by a diffeomorphism Φ_t for some $0\le t\le T$ (in other words, $\Phi_t(\gamma(u))=\gamma(v)$) and the homotopy class of the path $\xi_2^{-1}\circ\Phi_t(\gamma(u))\circ\xi_1$ is τ. Let us study separately each of the three parts of the boundary ∂C^ε. Denote by ∂_- the part of ∂C^ε on which $u=\varepsilon$, by ∂_+ the part on which $v=1-\varepsilon$, and $\partial_=$ the part on which $u+\varepsilon=v$ (with the induced orientation).

Denote by $\langle\gamma_1,\gamma_2\rangle^\tau$ the coefficient of τ in $\langle\gamma_1,\gamma_2\rangle$, see (3).

Lemma 14. *We have* $\mathrm{I}(\mathrm{ev}_\tau(\partial_-),W_T)=\langle\gamma,\Phi_-\rangle^\tau$ *and* $\mathrm{I}(\mathrm{ev}_\tau(\partial_+),W_T)=\langle\Phi_+,\gamma\rangle^\tau$. *Thus* $\sum_\tau \mathrm{I}(\mathrm{ev}_\tau(\partial_-),W_T)+\mathrm{I}(\mathrm{ev}_\tau(\partial_+),W_T)=2\,\mathrm{ind}_T(\gamma,p)$

Proof. For $(u,v)\in\partial_-$ we have $u=\varepsilon$ and $2\varepsilon\le v\le 1-\varepsilon$, thus we are interested in points of intersection of $\Phi_t(\gamma(\varepsilon))$ with $\gamma(v)$ for $0\le t\le T$ and $2\varepsilon\le v\le 1-\varepsilon$. But for small ε these are just intersections of Φ_+ (see Section 2.2) with the whole of γ. Moreover, it is easy to check that the signs of such intersection points coincide with the signs $\mathrm{sgn}(d)$, $d\in D(\Phi_+,\gamma)$ introduced in Section 2.1, which proves the first equality of the lemma. The second equality is proven in the same way, after we notice that $\varepsilon\le u\le 1-2\varepsilon$ and $v=1-\varepsilon$ correspond to intersections of $\Phi_t(\gamma(u))$ with $\gamma(v)=p$, i.e., of γ with Φ_-. □

Lemma 15. *We have* $\mathrm{I}(\mathrm{ev}_{[\gamma]}(\partial_=),W_T)=w(\gamma,X)$ *and* $\mathrm{I}(\mathrm{ev}_\tau(\partial_=),W_T)=0$ *for* $\tau\ne[\gamma]$.

Proof. For $(u,v)\in\partial_=$ we have $u+\varepsilon=v$. For small ε, values of u for which $\Phi_t(\gamma(u))=\gamma(u+\varepsilon)$ correspond to points in which γ' looks in the direction of X. Each such point gives the homotopy class $\tau=[\gamma]$ and is counted with a positive sign, if γ' turns counter-clockwise relative to X in a neighborhood of $\gamma(t)$ and with a negative sign otherwise. Comparison with the definition of $w(\gamma,X)$ in Section 1.1 proves the lemma. □

This concludes the proof of Theorem 5.

Finally, to prove Theorem 9 we need another configuration space. Call K^ε the set of pairs of points (u,v) on a circle such that the length of the arc $\overline{uv}$ and the

length of the arc $\overline{vu}$ are both $\geq \varepsilon$. Topologically, K^ε is an annulus, and its boundary $\partial\,\mathrm{K}^\varepsilon$ is a union of two circles L_+ and L_- defined by the conditions $|\overline{vu}| = \varepsilon$ and $|\overline{uv}| = \varepsilon$, respectively. The rest of the proof copies the proof of Theorem 5.

References

[1] Yu. Burman, M. Polyak, Geometry of Whitney-type formulas, *Moscow Math. Journal*, V.3 **3** (2003), pp. 823–832.

[2] M. Polyak, New Whitney-type formulae for plane curves, *AMS Transl.* **2** 190 (1999), pp. 103–111

[3] S. Smale, The classification of immersions of spheres in euclidean spaces. *Ann. Math.* (1959), pp. 327–344.

[4] V. Turaev, Intersections of loops in two-dimensional manifolds, *Math of the USSR–Sbornik* **35(2)** (1979), pp. 229–250.

[5] H. Whitney, On regular closed curves in the plane, *Compositio Math.*, **4** (1937), pp. 276–284.

121002, Independent University of Moscow, 11, B. Vlassievsky per., Moscow, Russia
E-mail address: burman@mccme.ru

Department of Mathematics, Technion- Israel Institute of Technology, Haifa 32000, Israel
E-mail address: polyak@math.technion.ac.il